

**RX SUMM—SYSTEMIC/SURGERY SEQ**

Item Length: 1  
 NAACCR Item #1639  
 Source of Standard: C0C  
 Rev (01/10, 01/11, 01/12, 06/12)  
 Dx Yr Req by MCR: 2006+

**Description:** *Records the sequencing of systemic therapy and surgical procedures given as part of the first course of treatment.*

**Instructions for Coding (See FORDS Revised for 2012, p. 277):**

- *Systemic/Surgery Sequence* is to be used for patients diagnosed on or after January 1, 2006.
- Code the administration of systemic therapy in sequence with the first surgery performed, described in the item *Date of First Surgical Procedure* (NAACCR Item #1200).
- If none of the following surgical procedures was performed: *Surgical Procedure of Primary Site* (NAACCR Item #1290), *Scope of Regional Lymph Node Surgery* (NAACCR Item #1292), *Surgical Procedure/Other Site* (NAACCR Item #1294), then this item should be coded 0.
- If the patient received both systemic therapy and any one or a combination of the following surgical procedures: *Surgical Procedure of Primary Site* (NAACCR Item #1290), *Scope of Regional Lymph Node Surgery* (NAACCR Item #1292), or *Surgical Procedure/Other Site* (NAACCR Item #1294), then code this item 2–9, as appropriate.
- If multiple first course treatment episodes were given such that both codes 4 and 7 seem to apply, use the code that defines the first sequence that applies.

Code	Label	Definition
0	No systemic therapy and/or surgical procedures	No systemic therapy was given; and/or no surgical procedure of primary site; no scope of regional lymph node surgery; no surgery to other regional site(s), distant site(s), or distant lymph node(s); or no reconstructive surgery was performed. It is unknown whether both surgery and systemic treatment were provided.
2	Systemic therapy before surgery	Systemic therapy was given before surgical procedure of primary site; scope of regional lymph node surgery; surgery to other regional site(s), distant site(s), or distant lymph node(s) was performed.
3	Systemic therapy after surgery	Systemic therapy was given after surgical procedure of primary site; scope of regional lymph node surgery; surgery to other regional site(s), distant site(s), or distant lymph node(s) was performed.

**Maine Cancer Registry Data Collection Manual for Hospitals, Fourth Edition - 2012**

<b>Code</b>	<b>Label</b>	<b>Definition</b>
4	Systemic therapy both before and after surgery	At least two courses of systemic therapy were given, at least one before and at least one more after a surgical procedure of primary site, scope of regional lymph node surgery, surgery to other regional site(s), distant site(s), or distant lymph node(s) was performed.
5	Intraoperative systemic therapy	Intraoperative systemic therapy was given during surgical procedure of primary site; scope of regional lymph node surgery; surgery to other regional site(s), distant site(s), or distant lymph node(s).
6	Intraoperative systemic therapy with other therapy administered before or after surgery	Intraoperative systemic therapy was given during surgical procedure of primary site; scope of regional lymph node surgery; surgery to other regional site(s), distant site(s), or distant lymph node(s) with other systemic therapy administered before or after surgical procedure of primary site; scope of regional lymph node surgery; surgery to other regional site(s), distant site(s), or distant lymph node(s) was performed.
7	Surgery both before and after systemic therapy	Systemic therapy was administered between two separate surgical procedures to the primary site; regional lymph nodes; surgery to other regional site(s), distant site(s), or distant lymph node(s).
9	Sequence unknown	Both surgery and systemic therapy were provided, but the sequence is unknown.